

Investigation of the dynamic behavior in Prussian blue analogs by picosecond time-resolved XAFS

Shunsuke Nozawa¹, Tokushi Sato¹, Kouhei Ichiyanagi², Ayana Tomita³, Hiroko Tkoro⁴, Shin-ichi Ohkoshi⁴, Shin-ichi Adachi¹, Shin-ya Koshihara³

¹Photon Factory, High Energy Accelerator Research Organization.

²Graduate School of Frontier Sciences, The University of Tokyo.

³Frontier Research Center & Department of Materials Science, Tokyo Institute of Technology.

⁴Department of Chemistry, School of Science, The University of Tokyo.

Prussian blue analogs have recently attracted great interest due to their various characteristics in the photo-induced phase transition (PIPT) as a photo-induced magnetization as well as photo-induced structural change [1-3]. The investigation of the dynamics of PIPT allows us to obtain information how the photo-excitation in the single site expands into a macroscopic phase transition. In this study, we have studied their dynamical behavior by means of the time-resolved x-ray absorption fine structure (XAFS) measurement. In time-resolved XAFS spectrum, extended x-ray absorption fine structure (EXAFS) is useful to reveal the dynamics of local structures around the absorbing atom. Moreover, the dynamics of electronic properties and the spin states are provided by x-ray absorption near edge structure (XANES).

The picosecond time-resolved XAFS experiments were performed on the NW14A at the Photon Factory Advanced Ring (PF-AR) [4]. Transient XAFS spectra were collected by pump-probe technique with the femtosecond laser system. Dynamical features induced by laser pulse excitation were obtained at 100 ps resolution. The detailed results will be presented.

Reference

- [1] H. Tokoro et al. *Appl. Phys. Lett.* **82** (2003) 1245.
- [2] Y. Sato et al. *J. Appl. Phys.*, **92** (2002) 4834.
- [3] T. Yamauchi et al., *Phys. Rev. B*, **72** (2005) 214425.
- [4] S. Nozawa et al. *J. Synchrotron Rad.*, **14** (2007) 313.